

Arysta LifeScience offers an expanded portfolio of innovative and trusted crop protection products for the pome fruit market — products that protect your apples and pears from weeds, insects and disease for improved fruit quality and yield.

Acramite*

ACRAMITE® provides good knockdown and long-lasting residual control of pest mites and is especially effective during the motile stage. ACRAMITE is also easy on predatory mites and beneficial insects. The unique mode of action makes it an excellent choice for a resistance management program.



CASORON® is an easy-to-use granular herbicide that provides long-lasting, effective control on a broad spectrum of annual grasses, broadleaf weeds and certain perennial weeds.

Kanemite^{*}

KANEMITE® provides excellent control of two-spotted and European red mite with a single rate of application. It is active on all life stages, providing quick knockdown of nymphs and adults while being easy on predatory mites and beneficial insects.



KASUMIN® is a novel bactericide. The active ingredient, kasugamycin, has a unique mode of action, making it effective against Fire blight as well as streptomycinresistant strains of bacteria in apples and pears.

Maestro

MAESTRO® provides dependable, broad-spectrum disease control in apples and pears with minimal impact on beneficial insects, making it a good fit for IPM programs. MAESTRO features multiple modes of action, which reduces the potential for disease resistance.



RIMON® controls codling moth and Oriental fruit moth in apples. The unique mode of action also makes it an excellent choice for a resistance management program.

Recommended Application Timing



	ACTIVE INGREDIENT	HERBICIDE / FRAC / IRAC GROUP	REI	PHI	RATE(S) / Ha	MAX NO. APPLICATIONS / SEASON	WEEDS / INSECTS / DISEASE CONTROLLED	NOTES
Lasoron	DICHLOBENIL	20	12 HOURS	N/A	110 - 175 KG	175 KG / HA / YEAR	ANNUAL GRASSES AND BROADLEAF / CERTAIN PERENNIAL WEEDS	APPLY TO BEARING OR ESTABLISHED SECOND-YEAR ORCHARD.
Maestro	CAPTAN	М	48 HOURS	7 DAYS	1.9 - 3.75 KG	N/A	APPLE SCAB, SOOTY BLOTCH, FLY SPECK, BROOK'S SPOT, BULL'S-EYE ROT, BITTER ROT, BLACK ROT	
Ka <u>su</u> min	KASUGAMYCIN	24	12 HOURS	90 DAYS	100 PPM (5.0 L / 1,000 L)	4 APPLICATIONS	FIRE BLIGHT	NOTE WATER VOLUME CHART ON THE LABEL TO HELP WITH RATE OF APPLICATION.
Rim <u>o</u> n	NOVALURON	15	12 HOURS	14 DAYS	0.93 - 1.4 L / 1,000 L	4 APPLICATIONS	CODLING MOTH, ORIENTAL FRUIT MOTH	MAY ALSO BE USED TO CONTROL APPLE CLEARWING MOTH AND DOGWOOD BORER AS A TRUNK SPRAY.
Kanemite	ACEQUINOCYL	20B	12 HOURS	14 DAYS	2.1 L	2 APPLICATIONS	TWO-SPOTTED SPIDER MITE, EUROPEAN RED MITE	THOROUGH COVERAGE IS KEY FOR CONTROL.
Acramite	BIFENAZATE	UN	12 HOURS	7 DAYS	568 - 851 GRAMS	1 APPLICATION	TWO-SPOTTED SPIDER MITE, EUROPEAN RED MITE	MOST EFFECTIVE DURING MOTILE STAGE OF MITES. USE HIGHER RATE FOR ERM CONTROL.

To learn more about the Arysta LifeScience portfolio of crop protection products, contact your crop protection retailer, visit www.gowithwhatworks.ca, or call 866-761-9397.

