

# Cabrio Plus fungicide

## technology sheet

Stop early and late blight in potatoes and get the proven benefits of **AgCelence**<sup>®</sup>.

- Two modes of action for control of early and late blight
- Unique **AgCelence** benefits<sup>1</sup> for disease control and better management of stress, for increased yield potential

### Active ingredients

Pyraclostrobin – Group 11  
Metiram – M<sup>2</sup>

### Formulation

Water dispersible granules

### One case contains

20 kg bag

### Storage

Store in cool, dry, ventilated area without floor drain.

### Crop

Potatoes

### Timing

7 to 14 day interval (for early blight)  
7 to 10 day interval (for late blight)



During periods of rapid growth or high disease pressure, use a shorter interval.

Make first application at tuber initiation before row closure.

Subsequent applications can be made:

- At row close to allow good coverage of the lower canopy where disease usually starts.
- Before harvest if the crop is under stress from heat/cold/drought conditions.

Cabrio<sup>®</sup> Plus fungicide provides a convenient formulation for easy mixing.



Source: BASF

### Diseases controlled

Early blight (*Alternaria solani*)

Late blight (*Phytophthora infestans*)

<sup>1</sup> **AgCelence** benefits refer to products that contain the active ingredient pyraclostrobin. All comparisons are against grower standard unless otherwise stated.

# Cabrio® Plus

Fungicide

## Application rates

**One jug of Cabrio Plus treats 8.9 ha (22 acres).**

Potatoes 2.25 kg/ha (0.91 kg/ac)

---

## Mixing order

1. Ensure the spray tank is clean before use.
  2. Fill the tank 1/2 full of water and start agitation.
  3. Add the required amount of Cabrio Plus fungicide to the tank and allow to completely dissolve before adding any tank-mix partners.
  4. Continue agitation while filling the remainder of the spray tank.
  5. After use, clean the spray tank according to label precautions.
- 

## Application tips

**Rainfastness** – 1 hour (pyraclostrobin). (Metiram will wash off in heavy rain)

**Restricted entry interval** – 12 hours.

**Resistance management** – Rotate to a non-Group 11 fungicide after each use. Do not exceed three (3) applications per year.

---

## Pre-harvest Interval

3 days for potatoes.

---

## Tank mixes

None on label.

Contact **AgSolutions**® Customer Care or your BASF Sales Representative for information on supported tank mixes.


---

## For more information

Call **AgSolutions** Customer Care at 1-877-371-BASF (2273) or visit [agsolutions.ca](http://agsolutions.ca).

---

### Average yield increase: Cabrio Plus in potatoes



In field-scale applications, Canadian growers saw an average increase of 32.9 cwt/acre over standard practice. Each bar represents 1 of 124 trials over 9 years. Source: Summary of trials in Canada 2006 to 2014

### Yield trial



Grower Standard

Cabrio Plus

Source: Burr Bros Farms, Delta, B.C. 2013

## Always read and follow label directions.

**AgSolutions** is a registered trade-mark of BASF Corporation; **AgCelence**, and **CABRIO** are registered trade-marks of BASF SE; all used with permission by BASF Canada Inc. . CABRIO PLUS fungicide should be used in a preventative disease control program. © 2015 BASF Canada Inc.