

GROUP

7

FUNGICIDE

APROVIA® Fungicide
SOLATENOL® technology

FUNGICIDE

EMULSIFIABLE CONCENTRATE


COMMERCIAL - AGRICULTURAL

For broad-spectrum disease control or suppression in labelled crops.

ACTIVE INGREDIENT:

Benzovindiflupyr..... 100 g/L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN



POISON
DANGER: CORROSIVE TO EYES & SKIN

REGISTRATION NO: **31981**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **0.5 – 1000 L**

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

Label

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically. Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

CORROSIVE to the eye. DO NOT get in eyes. Causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if inhaled. Avoid inhaling/breathing sprays. Severely irritating to the skin. DO NOT get on skin.

Restricted Entry Interval (REI): With the exception of grapes, **DO NOT** enter or allow worker entry into treated fields for 12 hours after application (12 hr REI). For cane turning and girdling in grapes, do not re-enter the treated area for 4 days. For all other activities in grapes, do not enter the treated areas for 12 hours.

Apply only when the potential for drift to areas of human habitation or activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

DO NOT apply by air.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield) during mixing, loading, application, clean-up and repair. Protective eyewear, chemical-resistant gloves and coveralls are not required during application within a closed cab and/or cockpit.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

Benzovindiflupyr is persistent and may carryover. It is recommended that any products containing benzovindiflupyr not be used in areas treated with this product during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

7.0 STORAGE

Store this product away from food or feed. Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area out of the reach of children and animals. This product does not require heated storage.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

GROUP	7	FUNGICIDE
-------	----------	-----------

APROVIA® Fungicide
SOLATENOL® technology

FUNGICIDE

EMULSIFIABLE CONCENTRATE


COMMERCIAL - AGRICULTURAL

For broad-spectrum disease control or suppression in labelled crops.

ACTIVE INGREDIENT:

Benzovindiflupyr..... 100 g/L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN



POISON
DANGER: CORROSIVE TO EYES & SKIN

REGISTRATION NO: **31981**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

Pamphlet

Table of Contents

	<u>Section Number</u>
NOTICE TO USER	1.0
FIRST AID	2.0
TOXICOLOGICAL INFORMATION	3.0
PRECAUTIONS	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE)	5.0
ENVIRONMENTAL PRECAUTIONS	6.0
STORAGE	7.0
DISPOSAL	8.0
PRODUCT INFORMATION	9.0
General Use Precautions	9.1
DIRECTIONS FOR USE.....	10.0
General Informaiton.....	10.1
Ground Application	10.2
Mixing Instructions	10.2.1
Spraying Instructions	10.2.2
Equipment Clean-Up	10.2.3
Airblast Application	10.3
Sprayng Instructions	10.3.1
Equipment Clean-Up	10.3.2
Product Specific Precautions.....	10.4
Spray Buffer Zones.....	10.5
Rotational Crop Restrictions	10.6
CROP USE DIRECTIONS.....	11.0
Potatoes – In Furrow Application	11.1
Fruiting Vegetables (Crop Group 8-09)	11.2
Cucurbit Vegetables (Crop Group 9)	11.3
Pome Fruit (Crop Group 11-09)	11.4
Blueberries (Lowbush).....	11.5
Small Fruit Vine Climbing Subgroup (Crop Subgroup 13-07F).....	11.6
MINOR USES	12.0
Bulb Onion (Subgroup 3-07A) and Green Onion (Subgroup 3-07B)	12.1
RESISTANCE MANAGEMENT RECOMMENDATIONS	13.0

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically. Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

CORROSIVE to the eye. DO NOT get in eyes. Causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if inhaled. Avoid inhaling/breathing sprays. Severely irritating to the skin. DO NOT get on skin.

Restricted Entry Interval (REI): With the exception of grapes, **DO NOT** enter or allow worker entry into treated fields for 12 hours after application (12 hr REI). For cane turning and girdling in grapes, do not re-enter the treated area for 4 days. For all other activities in grapes, do not enter the treated areas for 12 hours.

Apply only when the potential for drift to areas of human habitation or activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

DO NOT apply by air.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and protective eyewear (goggles or face shield) during mixing, loading, application, clean-up and repair. Protective eyewear, chemical-resistant gloves and coveralls are not required during application within a closed cab and/or cockpit.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

All users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

Benzovindiflupyr is persistent and may carryover. It is recommended that any products containing benzovindiflupyr not be used in areas treated with this product during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

7.0 STORAGE

Store this product away from food or feed. Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area out of the reach of children and animals. This product does not require heated storage.

8.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

For returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

9.0 PRODUCT INFORMATION

APROVIA® is a broad-spectrum product and is recommended for the control or suppression of many important plant diseases. APROVIA provides disease control of many rust, leaf spots and powdery mildews. APROVIA is primarily applied as a foliar spray and can be used in block, or alternating spray programs with other crop protection products. All applications should be made according to the use directions listed in the label.

9.1 General Use Precautions

Do not apply APROVIA through any type of irrigation equipment. **DO NOT** apply by air.

DO NOT USE IN GREENHOUSES (ALONE OR IN TANK MIXTURES)

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, AND/OR ILLEGAL RESIDUES.

Apply the higher rate and shorter interval under conditions of high disease pressures.

Unless indicated otherwise in the use directions for specific crops, make no more than two sequential applications of APROVIA before alternating with a non-Group 7 fungicide registered for the same disease.

Not all members of the crop groups have been tested for efficacy and phytotoxicity at the recommended label rates, and should be used at the discretion of the user.

10.0 DIRECTIONS FOR USE

10.1 General Information

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-877-SYNGENTA (1-877-964-3682) for information before applying any tank mix that is not specifically recommended on this label.

Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank-mixes under all environmental conditions or for all crop varieties. Tank mixes that are not specifically listed on this label should be tested on a small area first, under local conditions and using standard practices, to confirm the tank mix is suitable for widespread application.

To determine the physical compatibility of this product with other products, use a jar test.

Always read and follow label directions including WALES mixing order.

Thorough coverage is necessary to provide good disease control. Applications using nozzles, sufficient water volume and pressure to provide thorough and uniform coverage generally provide the most effective disease control.

To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap, as crop injury may occur.

For more information on spray equipment and calibration, consult sprayer manufacturers and provincial recommendations.

Do not apply through any ultra-low volume (ULV) spray system.

10.2 Ground Application

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural and Biological Engineers (ASABE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

10.2.1 Mixing Instructions

Prepare only the amount of spray mixture that is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Vigorous agitation is necessary for proper dispersal of the product. Maintain maximum agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

- APROVIA is an emulsifiable concentrate (EC) formulation.
- Prepare no more spray mixture than is required for the immediate operation.
- Thoroughly clean spray equipment before using this product.
- Agitate the spray solution before and during application.
- Rinse spray tank thoroughly with clean water after each day's use and dispose of pesticide rinsate by application to an already treated area.

APROVIA Alone (No Tank Mix)

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add APROVIA to the tank.
- Continue agitation while adding the remainder of the water.
- Begin application of the spray solution after APROVIA has completely dispersed into the mix water.
- Maintain agitation until all of the mixture has been sprayed.

APROVIA + Tank Mixtures: APROVIA is usually compatible with all tank-mix partners listed on this label. To determine the physical compatibility of APROVIA with other products, use a jar test. Using a jar, add the products to water in the same ratio and order as planned in the spray tank. Add wettable powders and water dispersible granular products first, then liquid flowables, then emulsifiable concentrates and solutions last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same order of addition required ingredients to the spray tank.

Mixing in the Spray Tank

- Add $\frac{1}{2}$ - $\frac{2}{3}$ of the required amount of water to the spray or mixing tank.
- With the agitator running, add the tank-mix partner(s) into the tank in the same order as described above.
- Allow the material to completely dissolve and disperse into the mix water. Continue agitation while adding the remainder of the water and APROVIA to the spray tank.
- Allow APROVIA to completely disperse.
- Spray the mixture with the agitator running.

10.2.2 Spraying Instructions

1. Water Volume: Apply in a minimum spray volume of 150 L/ha OR the volume given in the crop and pest specific instructions tabulated below, whichever is LARGER.
2. Sprayer Agitation: Use a jet agitator or liquid sparge tube which recirculates 7-10% of the tank per minute. **DO NOT** use an air sparger.
3. Pump: Screens should be used to protect the pump and prevent clogging. Use 16 mesh or *coarser* screens on the suction side of the pump. **DO NOT** place a screen in the recirculation line. Use 50 mesh or *coarser* screens between the pump and boom unless directed otherwise by the sprayer manufacturer.
4. Spray Nozzles: 80° or 110° drift reducing flat fan (e.g. those with a pre-orifice or turbulence chamber) or air induction nozzles are recommended. Use 50 mesh nozzle screens or as recommended by the nozzle manufacturer. **DO NOT** use flood type nozzles, controlled droplet application equipment, spray foils or hollow cone nozzles.
5. Pressure: As recommended by the nozzle manufacturer to achieve no smaller than ASABE fine sized droplets.
6. Apply at uniform speed and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to avoid potential crop injury from over application.

10.2.3 Equipment Clean-Up

Before Spraying:

- Prior to using APROVIA Fungicide, ensure that the spray tank, lines and filter are thoroughly clean.

After Spraying:

Thoroughly clean application equipment immediately after spraying. **DO NOT** allow APROVIA Fungicide residue to dry within the spray tank.

When using tank mixes, consult the tank-mix partner label for additional clean-up instructions. The following recommendations are provided:

1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **DO NOT** clean the sprayer near desirable vegetation, wells or other water sources.
2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial regulations.

10.3 Airblast Application

Airblast application: **DO NOT** apply during periods of dead calm. Avoid application of this product

when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

10.3.1 Spraying Instructions

1. Water Volume: Apply in a minimum spray volume of 375 L/ha OR the volume given in the crop and pest specific instructions tabulated below, whichever is LARGER. Water volume should exceed the minimum recommendation at later stages of fruit development.
2. Spray Quality: Select nozzles and pressure to achieve a minimum of ASABE fine sized droplets.
3. Spray Distribution: Select nozzles, orient deflectors, and adjust air speed and volume to ensure only the canopy is sprayed. Spray should just reach the top of the target. Account for the shape and canopy density of the target when setting spray distribution.
4. Apply at uniform speed and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to avoid potential crop injury from over application.

10.3.2 Equipment Clean-Up

Before Spraying:

Prior to using APROVIA, ensure that the spray tank, lines and filter are thoroughly clean.

After Spraying:

Thoroughly clean application equipment immediately after spraying. **DO NOT** allow APROVIA residue to dry within application equipment.

When using tank mixes, consult the tank-mix partner label for additional clean-up instructions. The following recommendations are provided:

1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **DO NOT** clean application equipment near desirable vegetation, wells or other water sources.
2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial regulations.

10.4 Product Specific Precautions

Follow recommendations provided by local disease monitoring services or provincial spray calendars regarding the appropriate application timing for protectant fungicides in your area. Cultural practices such as canopy management and removal of overwintered plant debris should be integrated with the use of fungicides to reduce disease incidence. Use the higher rate, and the shorter application interval under conditions of heavy infection pressure, on highly susceptible varieties, or when environmental conditions are favourable for disease development.

APROVIA treatments should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, plant residue management, crop rotation, and proper timing and placement of fertilizer and irrigation. Fungicide applications should begin prior to disease infection and continue throughout the season following a resistance management strategy (see **Resistance Management Recommendations**).

10.5 Spray Buffer Zones

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

Use of low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage, soil drench or soil incorporation do not require a spray buffer zone.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial habitat
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Potatoes, fruiting vegetables, cucurbit vegetables, onions		15	2	1	1	1
Airblast	Small fruit vine climbing subgroup	Early growth stage	50	25	15	5	3
		Late growth stage	40	15	5	3	2
	Pome fruit	Early growth stage	45	20	10	3	2
		Late growth stage	35	10	4	2	1

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the [Canada.ca](https://www.canada.ca) web site.

10.6 Rotational Crop Restrictions

Please see the following table for the crop rotational restrictions:

Rotational Crops	Planting Time From Last APROVIA Application
Potatoes and tuberous and corm vegetable subgroup (CG 1C) Dried shelled pea and bean subgroup (CG 6C) Soybeans Fruiting vegetables crop group (CG 8-09) Cucurbit vegetables crop group (CG 9) Cereals (wheat, barley, oats, rye, triticale) Corn (field, sweet, pop and specialty) Rapeseed subgroup (CG 20A) Onions (Crop sub-groups 3-07A and 3-07B)	0 days
All other crops Intended for Food and Feed	6 months/180 days

11.0 CROP USE DIRECTIONS

11.1 Potatoes – In Furrow Application

Crop	Potatoes – In Furrow Application
Rate	500-750 mL/ha See Rate Conversion Chart – Potatoes for rates per 100 m row
Diseases Suppressed	Stem and stolon canker and black scurf (<i>Rhizoctonia solani</i>)
Rate	750 mL/ha See Rate Conversion Chart – Potatoes for rates per 100 m row
Diseases Suppressed	Verticillium wilt (<i>Verticillium dahliae</i>)
Rate	500-750 mL/ha See Rate Conversion Chart – Potatoes for rates per 100 m row
Diseases Suppressed	Silver scurf (<i>Helminthosporium solani</i>)
Application Timing/Instructions	Make an in-furrow application in a water volume of 50-150 L per hectare at planting. Mount the spray nozzle so the spray is directed into the furrow as a 15-20 cm band just before the seed is covered. Apply the spray in a narrow band over the seed piece. Do not apply by air.
Specific Restrictions	Do not apply more than 100 g a.i./ha of foliar benzovindiflupyr-containing products on potato crops per season if an in-furrow application was made.

RATE CONVERSION CHART - POTATOES

Rate mL product /ha	Rate in mL product /100 m row for specified row spacing									
	Row Spacing (cm)									
	75	80	85	90	95	100	105	110	115	120
500	3.8	4.0	4.3	4.5	4.8	5.0	5.3	5.5	5.8	6.0
750	5.6	6.0	6.4	6.8	7.1	7.5	7.9	8.3	8.6	9.0

11.2 Fruiting Vegetables (Crop Group 8-09)

Crop	Fruiting Vegetables (Crop Group 8-09) Complete list of crops (CG 8-09): Eggplant; Groundcherry; Okra; Pepino; Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); Tomatillo; Tomato
Rate	500-750 mL/ha See Rate Conversion Chart – Potatoes for rates per 100 m row
Diseases Controlled	Early blight (<i>Alternaria solani</i>), Anthracnose (<i>Colletotrichum</i> spp.), Powdery mildew (<i>Oidiopsis sicula</i>), Septoria leaf spot (<i>Septoria lycopersici</i>)
Application Timing/Instructions	Begin applications prior to disease development and continue throughout the season on a 7 day interval, starting at first fruit set for all diseases except Early blight. For Early blight apply on a 7-14 day interval starting prior to disease establishment. If disease pressure is high, use the highest rate. For best results, sufficient water volume must be used to provide thorough coverage. Apply as a broadcast foliar spray in a minimum of 150 L/ha of water for thorough coverage.
Pre-Harvest Interval (PHI)	1 day
Maximum Amount of Product per Season	3 L/ha
Specific Restrictions	<ol style="list-style-type: none"> 1. Do not apply by air. 2. If used solo, apply APROVIA Fungicide in strict alternation with fungicides from a different cross-resistance group. 3. If use in mixture, do not apply more than 2 consecutive applications before switching to a non-Group 7 fungicide. Follow all precautions, restrictions and directions on the labels of fungicide products used in an alternation program.

11.3 Cucurbit Vegetables (Crop Group 9)

Crop	Cucurbit Vegetables (Crop Group 9) Complete list of crops (CG 9): Chinese waxgourd; Citron melon; Cucumber (field); Gerkin; Gourd, edible; Momordica spp.; Muskmelons (includes cantaloupe); Pumpkin; Squash, summer (includes zucchini); Squash, winter; Watermelon. Including cultivars and/or hybrids of these.
Rate	500-750 mL/ha
Diseases Controlled	Powdery mildew (<i>Sphaerotheca fuliginea</i> , <i>Erysiphe cichoracearum</i>) Alternaria leaf blight and spot (<i>Alternaria cucumerina</i> and <i>A. alternata</i>)
Rate	750 mL/ha
Diseases Controlled	Anthraxnose (<i>Colletotrichum orbiculare</i>), Cercospora leaf spot (<i>C. citrullina</i>), Gummy stem blight (<i>Didymella bryoniae</i>)
Application Timing/Instructions	Begin applications prior to disease development and continue throughout the season on a 7 day interval. If disease pressure is high, use the highest rate. For best results, sufficient water volume must be used to provide thorough coverage. Apply as a broadcast foliar spray in a minimum of 150 L/ha of water for thorough coverage. To control Gummy stem blight use 190 L/ha of water.
Pre-Harvest Interval (PHI)	1 day
Maximum Amount of Product per Season	3 L/ha
Specific Restrictions	<ol style="list-style-type: none"> 1. Do not apply by air. 2. Do not apply more than 2 consecutive applications before switching to a non-Group 7 fungicide. 3. Follow all precautions, restrictions and directions on the labels of fungicide products used in an alternation program.

11.4 Pome Fruit (Crop Group 11-09)

Crop	Pome Fruit (Crop Group 11-09) Complete list of crops: Apples, Crabapples, Pear, Pear (Asian), Quince, and Cultivars, varieties and/or hybrids of these commodities
Rate	300-500 mL/ha
Diseases Controlled	Scab (<i>Venturia inaequalis</i> and <i>V. pyrina</i>)
Rate	500 mL/ha
Diseases Controlled	Powdery mildew (<i>Podosphaera leucotricha</i>), Alternaria blotch (<i>Alternaria mali</i>)
Application Timing/Instructions	<p>Begin applications preventively. Apply APROVIA Fungicide alone or in combination with other non-Group 7 fungicides. For resistance management, apply no more than 2 consecutive sprays before alternating to a non-Group 7 containing fungicide. For best results, sufficient water volume must be used to provide thorough coverage. A minimum spray volume of 375 L/ha for ground applications is recommended.</p> <p>Protective Spray Schedule against Scab: Apply every 7-10 days starting at 6-13 mm (¼-½ inch) green tip or when environmental conditions become conducive for scab. If disease pressure is high, use the highest rate. Continue through petal fall until the threat of primary scab is complete. For improved fruit scab control, combine APROVIA Fungicide with a protectant fungicide registered to control scab (such as Allegro® 500F Fungicide, Dithane™ DG 75 Fungicide, Manzate® Pro-Stick™ Fungicide and Penncozeb® 75DF Fungicide). Only apply in combination on crops appearing on the labels of both partner products.</p> <p>Forecasting Spray Schedule against Scab: Use a forecasting system beginning at green tip. Apply within 48 hours of the onset of an infection period. Apply a follow-up spray after 7 days. For improved fruit scab control, combine APROVIA Fungicide with a protectant fungicide registered to control scab beginning at bloom (such as Allegro 500F, Dithane DG 75 Fungicide, Manzate Pro-Stick Fungicide and Penncozeb 75DF Fungicide). Only apply in combination on crops appearing on the labels of both partner products. For best results, sufficient water volume must be used to provide thorough coverage. A minimum spray volume of 375 L/ha for ground applications is recommended. The most restrictive directions must be followed.</p>
Pre-Harvest Interval (PHI)	30 days
Maximum Amount of Product per Season	2 L/ha
Specific Restrictions	<ol style="list-style-type: none"> 1. Do not apply by air. 2. Do not apply more than 2 consecutive applications before switching to a non-Group 7 fungicide. Follow all precautions, restrictions and directions on the labels of fungicide products used in an alternation program.

11.5 Blueberries (Lowbush)

Crop	Blueberries (Lowbush)
Rate	500-750 mL/ha
Diseases Controlled	Blueberry leaf rust (<i>Thekopsora minima</i>)
Rate	750 mL/ha
Diseases Suppressed	Valdensinia leaf spot (<i>Valdensinia heterodoxa</i>)
Application Timing/Instructions	Apply at the first sign of disease. A second application can be made 10-14 days after the first application, when disease pressure is severe or when agronomic or weather conditions are conducive to disease development or movement. A minimum spray volume of 200-300 L/ha for ground applications is recommended.
Pre-Harvest Interval (PHI)	365 days
Maximum Amount of Product per Season	1.5 L/ha
Specific Restrictions	<ol style="list-style-type: none"> 1. Do not apply by air. 2. Only apply in the non-cropping year of production (i.e. the vegetative or sprout phase of production). 3. Add non-ionic surfactant to achieve a final concentration of 0.2% v/v in the spray tank. If using an organo-silicone blend or vegetable based crop oil, the rate should be 0.125% v/v. 4. If used solo, apply APROVIA Fungicide in strict alternation with fungicides from a different cross-resistance group. 5. If used in mixture, apply SDHI fungicides in a maximum of 2 consecutive applications. 6. Crop safety has been found to be acceptable; however, not all possible blueberry clones have been tested under all conditions. It is recommended to test APROVIA on a representative portion of the crop and in local conditions to ensure that a phytotoxic response will not occur.

11.6 Small Fruit Vine Climbing Subgroup (Crop Subgroup 13-07F)

Crop	Small Fruit Vine Climbing Subgroup (Crop Subgroup 13-07F) Complete list of crops (CG 13-07F): Gooseberry, Grape, Maypop, Schisandra berry and cultivars, varieties and/or hybrids of these
Rate	500-750 mL/ha
Diseases Controlled	Powdery mildew (<i>Erysiphe</i> spp., <i>Sphaerotheca</i> spp.)
Application Timing/Instructions	For powdery mildew, begin at bud break and apply on a 7-21 day interval. Use the high rate and shorter interval when conditions are more conducive to disease. Apply in sufficient water volume to obtain thorough coverage. A minimum spray volume of 150 L/ha is recommended.
Pre-Harvest Interval (PHI)	21 days
Maximum Amount of Product per Season	3 L/ha
Specific Restrictions	<ol style="list-style-type: none"> 1. Do not apply by air. 2. If used solo, apply APROVIA Fungicide in strict alternation with fungicides from a different cross-resistance group. 3. If used in mixture, do not apply more than 2 consecutive applications before switching to a non-Group 7 fungicide. Follow all precautions, restrictions and directions on the labels of fungicide products used in an alternation program. 4. Do not apply more than three total applications of any Group 7 product per season on grapes. A fourth application is allowed where 12 or more fungicide applications are made.

12.0 MINOR USES

READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Syngenta Canada Inc. under the User Requested Minor Use Label Expansion program. For these uses, Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

12.1 Bulb Onion (Subgroup 3-07A) and Green Onion (Subgroup 3-07B)

Crops	Bulb onion (Subgroup 3-07A) and Green Onion (Subgroup 3-07B) Complete list of crops: Dry bulb: Onion, bulb; Onion, Chinese, bulb; Garlic, bulb; Garlic, great-headed (elephant); Shallot, bulb Green Onion: Green onions; Leek; Welch onion; Chives, fresh leaves; Chives, Chinese; Shallot, fresh leaves; Lady's leek, Onion, fresh' Onion, tree, tops; Wild leek
Disease Controlled	Purple blotch (<i>Alternaria porri</i>) and garlic rust (<i>Puccinia allii</i>)
Disease Suppressed	Leaf blight (<i>Stemphylium vesicarium</i>)
Use rate mL/ha	750 mL/ha
Application Timing/Instructions	Begin applications when the plants are 5-10 centimeters high or when conditions become favourable for disease development. Make applications on a 7 – 14 days interval. Use the shorter interval when conditions are more conducive to disease. Apply specified rates in a minimum of 150 L of water per hectare up to 600 L of water per hectare.
Pre-Harvest Interval (PHI)	7 days
Maximum Number of Applications per Season	4
Maximum Amount of Product per Season	3 L/ha
Specific Use Restrictions: <ol style="list-style-type: none"> 1. Do not apply by air. 2. Do not apply more than 2 consecutive applications before switching to a non-Group 7 fungicide. Follow all precautions, restrictions and directions on the labels of fungicide products used in an alternation program. 	

13.0 RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, APROVIA contains a Group 7 fungicide in the succinyl dehydrogenase inhibitor (SDHI) class. Any fungal population may contain individuals naturally resistant to benzovindiflupyr and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of benzovindiflupyr or other Group 7 fungicides with different groups that control the same pathogens.

Apply no more than 2 sequential applications unless otherwise stated in the crop section.

DO NOT apply at rates lower than recommended on the label.

Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant

resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for sign of resistance development. Notify Syngenta Canada Inc. if reduced sensitivity of the pathogen to APROVIA Fungicide is suspected.

If disease continues to progress after treatment with this product, DO NOT increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, to which pathogen resistance has not developed.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information or to report suspected resistance, contact company representatives at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

Application Limitations and Pre-harvest Intervals

Crop*	Maximum Product Rate/ha/application (mL/ha)	Maximum total (L/ha/Season)	Pre-Harvest Interval (PHI) (days)	Minimum Re-treatment Interval (days)
Potatoes	750 (soil)	0.75 (soil)	NA	NA
Fruiting Vegetables (CG 8-09)	750	3	1	7
Cucurbits (CG 9)	750	3	1	7
Pome (CG 11-09)	500	2	30	7
Blueberries	750	1.5	365	10
Small fruit and vine climbing subgroup (13-07F)	750	3	21	7
Onions (CG 3-07A and 3-07B)	750	3	7	7

*For specific crops in a group and use directions, refer to the Specific Directions For Use.

APROVIA® and SOLATENOL® are trademarks of a Syngenta Group Company.

Allegro®, Dithane™, Manzate®, Pencozeb® and Pro-Stick™ are trademarks of their respective companies.